arXiv:1604.01843v1 [math-ph] 7 Apr 2016

Renormalization group flow, Entropy and Eigenvalues

Dan Li *

Abstract

The irreversibility of the renormalization group flow is conjectured to be closely related to the concept of entropy. In this paper, the variation of eigenvalues under the renormalization group flow will be studied. Based on the one-loop approximation to the effective field theory, we will use the heat kernel method and zeta function regularization. In even dimensions, the variation of eigenvalues is given by the top heat kernel coefficient, and the conformal anomaly is relevant. In odd dimensions, we will conjecture a formula for the variation of eigenvalues based on the holographic principle.

1 Introduction

The idea of renormalization group (RG) emerges from the renormalization process of a quantum field theory (QFT) by which, if possible, all divergent infinities are taken care of and the QFT is made into a renormalizable theory. As a prototypical example, when the infinity in the momentum space is cut off by a large energy scale Λ, the coupling constants are now depending on the introduced renormalization parameter Λ and the original QFT becomes an effective field theory. In other words, one obtains different effective field theories parametrized by the cut-off scale in the space of quantum field theories (i.e., theory space).

As the energy scale varies, for example from the ultraviolet (UV) regime to the infrared (IR) regime, the renormalization group forms a renormalization group flow, which could have several fixed points. One basic assumption

*E-mail: li1863@math.purdue.edu

of the RG flow is that if the scale change is small from high energy to low energy, for example infinitesimal, i.e., $\Lambda \to \Lambda' = \Lambda - \delta\Lambda$, then the corresponding effective field theories are scale invariant. In most cases, scale invariance implies conformal invariance, so it has important applications in conformal geometry if the RG flow is approximated by some geometric flow. Of course, the fixed points, i.e., some conformal field theories (CFTs) in the theory space, play important roles in the study of RG flow. Another important observation of the RG flow is its irreversibility, which is conjectured to be intimately related to the concept of entropy. For exmaple, for any renormalizable 2d field theory, the Zamolodchikov c-theorem [16] tells us that the central charge at the fixed points of the RG flow decreases monotonically, i.e., $c_{UV} > c_{IR}$.

The beta function of a coupling parameter $g(\Lambda)$ depending on the energy scale Λ is defined as

$$\beta(g) := \Lambda \frac{\partial g}{\partial \Lambda} = \frac{\partial g}{\partial \ln \Lambda}$$

which describes the running of the coupling parameter g along with the renormalization scale Λ. At a fixed point of the RG flow, the conformal invariance implies the vanishing of the beta function, i.e., $\beta(g) = 0$. In metric geometry, the coupling parameter g is always linked to the metric g_{ij} of a Riemannian manifold.

Friedan first considered the perturbative RG flow of a two dimensional nonlinear sigma model, and computed the beta function of the metric g_{ij} of the target space based on Feynman diagram expansion [5, 6]. The one-loop approximation of the RG flow, i.e., the first term in the beta function, gives rise to the Ricci flow,

$$\frac{\partial g_{ij}}{\partial t} = -2R_{ij} \tag{1}$$

where $t = \ln(\Lambda^{-1})$ is the "time" parameter and R_{ij} is the Ricci curvature. As a remark, the parameter t has nothing to do with the physical time. The Ricci flow equation (1), which was first introduced by Hamilton, is always viewed as the geometric analogy of the diffusive heat equation. More recently, Perelman introduced the W-functional under the Ricci flow and proved the no local collapsing theorem in the resolution of the Poincaré conjecture and Thurston's geometrization conjecture [14]. As pointed out by Perelman in [14], the definition of W-functional was inspired by the concept of entropy in statistical mechanics, which is related to the RG flow heuristically.

The variation of eigenvalues of geometric operators such as the Laplacian under the Ricci flow has been studied in the literature, see [1, 4, 10, 13], and the monotonicity of eigenvalues has been established. The idea behind such monotonicity is that from the entropy functional such as the F-functional or W-functional by Perelman, eigenvalues of the corresponding geometric operators inherit the monotonicity from the entropy functional. And such entropy functional originates from the thermal entropy in statistical mechanics, for example see [11].

In this paper, we will consider the variation of eigenvalues under the RG flow, and express it in terms of spectral invariants, i.e., heat kernel coefficients and spectral zeta function. In mathematical physics, the top heat kernel coefficient plays an important role, for example, in explanations of the conformal anomaly, the chiral anomaly and the Atiyah–Singer index theorem [15]. It is not a surprise that the variation of eigenvalues equals the top heat coefficient a_{2k} in even dimensions. In general, there is no monotonicity of eigenvalues or equivalently positivity of a_{2k}, since a_{2k} consists of different parts expressed in scalar curvature, Ricci curvature tensor and Riemann curvature tensor. But it is possible to extract a positive or negative part such as the Euler density, the Weyl tensor or the Q-curvature so that the variation of eigenvalues has a positive or negative sign under some mild conditions on the Riemannian manifold. As a result, the regularized log-determinant has a term from the variation of eigenvalues. Of course, the variation of log-determinants can be computed by Polyakov formulas in conformal geometry.

In odd dimensions, the situation is more subtle. We know that the heat kernel coefficients $a_{2k-1} = 0$, and there is no conformal anomaly in odd dimensional CFTs. From classical statistical mechanics, we learned that the partition function and the free energy can be used interchangeably to express the probability function. Based on the holographic renormalization group flow, if the even dimensional bulk theory is characterized by the partition function, then the boundary theory should be described by the free energy. In addition, the irreversibility of the RG flow incarnates itself in a so-called C-function, which spiritually reflects the increasing of entropy. More precisely, there is a c-theorem in 2d [16], and there is an a-theorem in 4d [9], where a and c are central charges in CFTs and both theorems are closely related to the conformal anomaly. In 3d, there is an F-theorem [3], where F is the free energy, and it is also viewed as an entanglement entropy. So based on the holographic principle, we conjecture a formula for the variation of eigenvalues under the RG flow in odd dimensions.

This paper is organized as follows. In section 2, we will review some basic facts about the Polyakov action, perturbative RG flow and the heat kernel method. The non-perturbative heat kernel approach will recover the first order approximation to the RG flow, i.e., the Ricci flow. In section 3, we will derive the variation of eigenvalues under the RG flow in even dimensions, and the result is given by the top heat kernel coefficient. In section 4, we will review the thermal entropy and the irreversibility of the RG flow, then a formula for the variation of eigenvalues will be conjectured based on the holographic principle.

2 Polyakov action

In this section, we first review some facts about the perturbative renormalization group flow and the heat kernel method. In the last subsection, we look into the Laplacian operator from the Polyakov action and recover the Ricci flow from a heat kernel coefficient.

2.1 Perturbative RG flow

In order to study the geometry of a string propagating in a curved background, one considers the nonlinear sigma model (or σ-model) defined by the Polyakov action, which is the kinetic energy of a scalar field mapping a 2d surface Σ into the target space M. More precisely, for a smooth embedding $\phi : (\Sigma, \gamma) \to (M, g)$, the Polyakov action is defined as

$$S(g;\phi) = \frac{T}{2}\int_{\Sigma} dv\ \gamma^{\mu\nu}\partial_\mu\phi^i(x)\partial_\nu\phi^j(x)g_{ij}(\phi(x)) \tag{2}$$

where $dv = \sqrt{|\gamma|}d\sigma$ is the standard volume measure on Σ. In string theory, the Polyakov action describes a bosonic string model and the parameter T is the string tension. Basically, the Polyakov action computes the area of the string worldsheet in the curved spacetime. In the literature, the Regge slope α' is also widely used, and it is connected to T by $T = 1/2\pi\alpha'$. A general nonlinear sigma model contains dilatons and B-fields etc., in this section we focus on the Polyakov action (2), i.e., the free field theory.

For simplicity, we assume that the smooth Riemann surface (Σ, γ) and the n-dimensional ($n \geq 2$) Riemannian manifold (M, g) are oriented and compact without boundary. The space of smooth maps is denoted by $C^\infty(\Sigma, M)$, and

the scalar fields $\phi : \Sigma \to M$ are assumed to be smooth. The Polyakov action $S(g;\phi)$ is also assumed to be invariant under the diffeomorphism group of M, i.e., $Diff(M)$. In addition, we only consider Euclidean quantum field theory (QFT) in this paper.

The renormalization group (RG) flow of the nonlinear sigma model (2) was first investigated by Friedan [5, 6], an approach based on dimensional regularization of Feynman diagram evaluation can be found in [7], and the geometric picture of the RG flow is explained in [2].

We briefly recall the idea of perturbative RG flow based on regularization and renormalization following [2]. First one considers the effective action W defined by

$$e^{-W} = \int [D\phi] e^{-S(g;\phi)}$$

where the path integral measure is formally defined over scalar fields modulo diffeomorphic equivalence. When the parameter $T \to \infty$ or $\alpha' \to 0$, the effective action is roughly the classical action plus quantum fluctuations around the stationary points of the action $S(g;\phi)$. If there exists a decomposition $\phi = \phi_0 + \pi$ around the background field ϕ_0 such that $\frac{\delta S}{\delta \phi}(\phi_0) = 0$ and $\pi \in C^\infty(\Sigma, \phi^* TM)$ is a quantum fluctuation, then under the saddle point approximation the local contribution to W around ϕ_0 is approximated by

$$W(\phi_0) = S(\phi_0) - ln \int [D\pi] exp\left(-\frac{1}{2} S''_{\mu\nu}(\phi_0)\pi^\mu \pi^\nu\right)$$

Using the normal geodesic coordinates around ϕ_0, $S''_{\mu\nu}(\phi_0)$ is expanded out and the Riemann curvature R_{ijkl} appears naturally in the covariant derivatives. Applying the Feynman rules to one-loop Feynman diagrams, a momentum cut-off Λ is introduced into the Feynman propagator. After this regularization, the divergence part of the field theory only lies in a term with the logarithmic energy $\ln \Lambda$ multiplied by the Ricci curvature R_{ij}. Finally, the renormalization procedure identifies the Ricci curvature as the lowest order term in the beta function. More precisely, the first order term in α' in the RG flow equation gives rise to the Ricci flow,

$$\frac{dg_{ij}}{dt} = -2\alpha' R_{ij} + O(\alpha'^2)$$

where $t = \ln \Lambda^{-1}$ or $\Lambda = e^{-t}$.

The higher order terms in the RG flow can be obtained similarly by evaluating Feynman diagrams with more loops, and the second term was computed in Friedan's thesis [6],

$$\frac{dg_{ij}}{dt} = -\beta(g_{ij}) = -2\alpha' R_{ij} - \alpha'^2 R_i^{klm} R_{jklm} + O(\alpha'^3)$$

The perturbative RG flow is conjectured to be a good approximation to the nonperturbative RG flow. In the next section, we will look at a nonperturbative approach based on the heat kernel method. Because of the feature of heat kernel expansion, this approach only determines the one-loop approximation to the effective field theory.

2.2 Heat kernel expansion

Suppose (M, g) is a compact smooth Riemannian manifold without boundary, let Δ be the Laplace–Beltrami operator with eigenvalues

$$0 \leq \lambda_0 < \lambda_1 \leq \lambda_2 \leq \cdots \lambda_k \leq \cdots$$

If the corresponding eigenfunctions are given by $\{\psi_k\}$, which form an orthonormal basis for $L^2(M)$, then the heat kernel $K(t; x, y; \Delta)$ is defined as

$$K(t; x, y; \Delta) = \sum_k e^{-\lambda_k t} \psi_k(x) \psi_k(y)$$

The trace of the heat kernel, sometimes called the partition function $Z(t)$, is defined by

$$Z(t) = Tr(e^{-t\Delta}) = \sum_k e^{-t\lambda_k} = \int_M dV\, K(t; x, x; \Delta) = K(t, \Delta) \qquad (3)$$

which has an asymptotic expansion when t tends to zero,

$$Tr(e^{-t\Delta}) \sim (4\pi t)^{-\frac{n}{2}} (a_0 + a_2 t + a_4 t^2 + \cdots) \quad \text{as} \quad t \to 0^+$$

where $\dim M = n$ and the heat kernel coefficients a_{2k} are integrals of local spectral invariants, for example,

$$\begin{aligned} a_0 &= Vol(M, g) \\ a_2 &= \tfrac{1}{6} \textstyle\int_M dV\, R \\ a_4 &= \tfrac{1}{360} \textstyle\int_M dV\, (-12\Delta R + 5R^2 - 2R^{ij} R_{ij} + 2R^{ijkl} R_{ijkl}) \end{aligned}$$

One defines the spectral zeta function of the Laplace operator Δ as usual,

$$\zeta_\Delta(s) = \sum_{\lambda_k \neq 0} \lambda_k^{-s} = Tr(\Delta^{-s})$$

which is well-defined for $Re(s) > \frac{n}{2}$. Let $\Gamma(s)$ be the Gamma function, one has the regularized zeta function by the Mellin transform,

$$\zeta_\Delta(s) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1}(Tr(e^{-t\Delta}) - 1)dt$$

From the basic formula for a positive $\lambda > 0$,

$$\ln \lambda = -\int_0^\infty \frac{dt}{t} e^{-t\lambda}$$

it is easy to see that

$$\ln \det(\Delta) = Tr \ln(\Delta) = -\int_0^\infty \frac{dt}{t} Tr(e^{-t\Delta})$$

so the one-loop effective action is

$$W := \frac{1}{2} \ln \det(\Delta) = -\frac{1}{2} \int_0^\infty \frac{dt}{t} Tr(e^{-t\Delta})$$

In addition, the logarithm of the determinant (or log-determinant) of the Laplace operator is always expressed in terms of the regularized zeta function,

$$\ln \det \Delta = -\frac{d}{ds}\zeta_\Delta(s)|_{s=0} = -\zeta'_\Delta(0)$$

For a general Laplace-type operator $D = -\Delta + V$, where V is a matrix-valued potential, and an auxiliary smooth matrix-valued function f, the trace of the heat kernel is well defined,

$$Tr(fe^{-tD}) = \int_M dV\, K(t; x, x; D) f(x) = K(t, f, D)$$

There is an asymptotic expansion when t tends to 0,

$$Tr(fe^{-tD}) \sim (4\pi t)^{-\frac{n}{2}} [a_0(f, D) + a_2(f, D)t + a_4(f, D)t^2 + \cdots] \quad \text{as} \quad t \to 0^+$$

For example, the first three non-zero terms are given by

$$\begin{aligned} a_0(f,D) &= \int_M dV\ Tr\{f\} \\ a_2(f,D) &= \tfrac{1}{6}\int_M dV\, Tr\{f(6V+R)\} \\ a_4(f,D) &= \tfrac{1}{360}\int_M dV\, Tr\{f(60\Delta V + 60RV + 180V^2 \\ &\qquad +12\Delta R + 5R^2 - 2R^{ij}R_{ij} + 2R^{ijkl}R_{ijkl})\} \end{aligned}$$

Similarly, the spectral zeta function can be generalized as,

$$\zeta(s,f,D) = Tr(fD^{-s})$$

and by the Mellin transform,

$$\zeta(s,f,D) = \frac{1}{\Gamma(s)}\int_0^\infty dt\, t^{s-1} Tr(fe^{-tD})$$

The heat kernel coefficients can be expressed as the residues at the poles,

$$a_k(f,D) = Res_{s=(n-k)/2}(\Gamma(s)\zeta(s,f,D))$$

in particular,

$$a_n(f,D) = \zeta(0,f,D)$$

2.3 Ricci flow

The Polyakov action (2) is also written as

$$S(g;\phi) = \int_\Sigma dv\, tr_\gamma(\phi^* g)$$

where the string tension T has been absorbed into the action, and the Lagrangian is the trace of the pullback or induced metric,

$$\phi^* g_{\mu\nu}(x) = g_{ij}(\phi(x))\partial_\mu\phi^i(x)\partial_\nu\phi^j(x)$$

In particular, if the spacetime metric g is assumed to be flat, i.e., $g_{ij} = \delta_{ij}$, then one obtains the simplified action,

$$S(\delta;\phi) = \int_\Sigma d\sigma\, \sqrt{|\gamma|}\gamma^{\mu\nu}\partial_\mu\phi^i(x)\partial_\nu\phi_i(x)$$

If one further applies the integration by parts, then the action becomes

$$S(\delta;\phi) = -\int_\Sigma dv\, \phi^i(x)\Delta_\gamma\phi_i(x)$$

where Δ_γ is the Laplace–Beltrami operator with respect to $\gamma_{\mu\nu}$,

$$\Delta_\gamma u = \frac{1}{\sqrt{|\gamma|}}\partial_\mu(\sqrt{|\gamma|}\gamma^{\mu\nu}\partial_\nu u)$$

In practice, the spacetime metric is sometimes decomposed as $g_{ij} = \delta_{ij} + h_{ij}$, in this case the Polyakov action is a sum,

$$S(g;\phi) = S(\delta;\phi) + S(h;\phi)$$

where $S(h;\phi)$ is viewed as a fluctuation.

Lemma 1. *The Polyakov action* (2) *can be expressed as*

$$S(g;\phi) = -\int_\Sigma dv\, g_{ij}\phi^i\partial^\mu\partial_\mu\phi^j + \frac{1}{2}\int_\Sigma dv\,(\partial^\mu\partial_\mu g_{ij})\phi^i\phi^j \tag{4}$$

if both the worldsheet and spacetime metrics are symmetric.

Proof. First of all, it is natural to assume that the worldsheet metric $\gamma_{\mu\nu}$ is symmetric, i.e., $\gamma_{\mu\nu} = \gamma_{\nu\mu}$. The Polyakov action is independent of the metric γ of Σ, in practice γ is always assumed to be flat $\gamma_{\mu\nu} = \delta_{\mu\nu}$. The main tool in computing the following integrals is the integration by parts.

$$\begin{aligned} S(g;\phi) &= \textstyle\int_\Sigma dv\, g_{ij}(\phi)\partial^\mu\phi^i\partial_\mu\phi^j \\ &= -\textstyle\int_\Sigma dv\, \phi^i\,\partial^\mu[g_{ij}(\phi)\partial_\mu\phi^j] \\ &= -\textstyle\int_\Sigma dv\,[\phi^i g_{ij}(\phi)\partial^\mu\partial_\mu\phi^j + \phi^i\partial^\mu g_{ij}(\phi)\partial_\mu\phi^j] \end{aligned}$$

where the modified Laplace operator will be denoted by $\Delta_g = g_{ij}(\phi)\partial^\mu\partial_\mu$. If we further assume the spacetime metric g is symmetric, i.e., $g_{ij} = g_{ji}$, then the modified Laplace operator is formally self-adjoint, i.e., $\Delta_g^* = \Delta_g$.

Let us continue to work on the second integral,

$$\begin{aligned} &\textstyle\int_\Sigma dv\, \phi^i\partial^\mu g_{ij}(\phi)\partial_\mu\phi^j \\ &= -\textstyle\int_\Sigma dv\, \phi^j\partial_\mu[\phi^i\partial^\mu g_{ij}(\phi)] \\ &= -\textstyle\int_\Sigma dv\,[\phi^j\partial_\mu\phi^i\partial^\mu g_{ij}(\phi) + \phi^j\phi^i\partial_\mu\partial^\mu g_{ij}(\phi)] \\ &= \textstyle\int_\Sigma dv\, g_{ij}(\phi)\partial^\mu[\phi^j\partial_\mu\phi^i] - \int_\Sigma dv\,\phi^i\phi^j\partial_\mu\partial^\mu g_{ij}(\phi) \\ &= \textstyle\int_\Sigma dv\,[g_{ij}(\phi)\partial_\mu\phi^i\partial^\mu\phi^j + g_{ij}(\phi)\phi^j\partial^\mu\partial_\mu\phi^i] - \int_\Sigma dv\,\phi^i\phi^j\partial_\mu\partial^\mu g_{ij}(\phi) \end{aligned}$$

Since γ is symmetric, one has $S(g;\phi) = \int_\Sigma dv g_{ij}(\phi)\partial_\mu\phi^i\partial^\mu\phi^j$ as well. Putting together, we obtain

$$\begin{aligned} S(g;\phi) = \quad & -\int_\Sigma dv\, \phi^i\Delta_g\phi^j - S(g;\phi) \\ & -\int_\Sigma dv\, \phi^j\Delta_g\phi^i + \int_\Sigma dv\, \phi^i\phi^j\partial_\mu\partial^\mu g_{ij}(\phi) \end{aligned}$$

Or equivalently,

$$S(g;\phi) \quad = -\int_\Sigma dv\, \phi^i\Delta_g\phi^j + \tfrac{1}{2}\int_\Sigma dv\, \phi^i\Delta g_{ij}(\phi)\phi^j$$

where $\Delta = \gamma^{\mu\nu}\partial_\nu\partial_\mu = \partial^\mu\partial_\mu = \partial_\mu\partial^\mu$. If the spacetime metric g is flat, i.e., $g_{ij} = \delta_{ij}$, then the 2nd term is zero and the 1st term recovers the Laplace–Beltrami operator.

□

If the real scalar field $\phi : \Sigma \to M$ has the local coordinates $\phi(x) = (\phi^1(x), \cdots, \phi^n(x))$ for $x \in \Sigma$, then we can denote the above integrals by inner products,

$$\langle\phi, \Delta\phi\rangle_g := \int_\Sigma dv\, g_{ij}\phi^i\partial^\mu\partial_\mu\phi^j$$

$$\langle\phi, \phi\rangle_{\Delta g} := \int_\Sigma dv\, (\partial^\mu\partial_\mu g_{ij})\phi^i\phi^j$$

Similarly, the Polyakov action can be expressed as

$$S(g;\phi) = \int_\Sigma dv\, g_{ij}\partial^\mu\phi^i\partial_\mu\phi^j = \langle d\phi, d\phi\rangle_g$$

Lemma 2. *The relation (4) in the above lemma is equivalent to the following,*

$$\langle d\phi, d\phi\rangle_g = -\langle\phi, \Delta\phi\rangle_g + \frac{1}{2}\langle\phi, \phi\rangle_{\Delta g}$$

or under the standard inner product $\langle\phi, \phi\rangle := \sum_{i,j=1}^n \int_\Sigma dv\, \phi^i\phi^j$,

$$\langle d\phi, gd\phi\rangle = -\langle\phi, g\Delta\phi\rangle + \frac{1}{2}\langle\phi, \Delta g\phi\rangle$$

Therefore, the operator in the Polyakov action is given by

$$L(\phi, g, \gamma) = -g\Delta + \frac{1}{2}\Delta g = -g_{ij}(\phi)\gamma^{\mu\nu}\partial_\mu\partial_\nu + \frac{1}{2}\gamma^{\mu\nu}\partial_\mu\partial_\nu g_{ij}(\phi(x)) \qquad (5)$$

If we view g_{ij} as an auxiliary functioin, then in the standard form

$$L(\phi, g, \gamma) = g(-\Delta + V) = gD(\phi, g, \gamma)$$

where the potential $V = \frac{1}{2}g^{-1}\Delta g$ and $D = -\Delta + V$ is a Laplace-type operator. For example, if $g_{ij}(\phi) = e^{\phi(x)}g^0_{ij}$ for some constant metric g^0 in the background, then

$$D = -\Delta + \frac{1}{2}(\partial^\mu\phi\partial_\mu\phi + \Delta\phi)$$

If we apply the heat kernel method to the operator D and the auxiliary function g, then we obtain the partition function depending on g,

$$Z(t, g) = Tr(ge^{-tD})$$

and the one-loop effective action of the Polyakov action,

$$W(g) = \frac{1}{2}\ln\det(gD) = -\frac{1}{2}\int_0^\infty \frac{dt}{t}Tr(ge^{-tD})$$

Suppose the coupling parameter g is a function of the energy scale Λ, then the Callan–Symanzik equation reads,

$$\frac{\partial W(g(\Lambda))}{\partial \ln\Lambda} + \beta(g)\frac{\partial W(g)}{\partial g} = 0$$

So the beta function can be computed by,

$$\beta(g) = -\frac{\partial W(g(\Lambda))}{\partial \ln\Lambda}\Big/\frac{\partial W(g)}{\partial g}$$

If we approximate the denominator by

$$\frac{\partial W(g)}{\partial g} \sim -\frac{1}{2}\int_0^\infty \frac{dt}{t}Tr(e^{-tD}) = \frac{1}{2}\ln\det(D)$$

then it can be viewed as a normalization constant N.

From the next section, the regularized effective action will be

$$W(g(\Lambda)) = -\frac{1}{2}[\zeta(0, D) + 2\ln\Lambda\, a_n(g, D)]$$

In order to approximate the first order term in the beta function, it is enough to take $a_2(g, D)$, so we have

$$-\beta(g) = \frac{1}{N}\frac{\partial W(g(\Lambda))}{\partial \ln \Lambda} \sim -2a_2(g, D) = -\frac{1}{3}\int_M dV\, Tr(gR) - \int_M dV\, Tr(\Delta g)$$

If we view the first integral in $a_2(g, D)$ as a finte part, and the local form of the second integral gives rise to the Ricci flow,

$$\frac{\partial g_{ij}}{\partial t} = -2R_{ij} \sim \Delta g_{ij}$$

since one has the approximation $\Delta g_{ij} \sim -2R_{ij}$ in harmonic local coordinates.

3 Variation of eigenvalues in even dimensions

In this section, we will continue to consider the Polyakov action (2), and derive a formula for the variation of eigenvalues under the renormalization group (RG) flow.

Theorem 1. *The nonperturbative RG flow equation of the eigenvalues of the Laplacian operator in the Polyakov action is given by*

$$\frac{\partial \lambda}{\partial t} = a_n(g, D) = \zeta(0, g, D) \tag{6}$$

Proof. For a scalar field with upper energy bound Λ, i.e., $\phi \in C^\infty(\Sigma, M)[0, \Lambda]$, the effective action $S_{eff}[\Lambda, g]$ is defined by

$$e^{-S_{eff}[\Lambda,g]} = \int_{C^\infty(\Sigma,M)[0,\Lambda]} D\phi\ exp\left\{-\langle\phi, L(\Lambda, \phi, g)\phi\rangle\right\}$$

where $L(\Lambda, \phi, g) = L(\phi, g, \gamma)$ from (5) is the Laplacian operator in the Polyakov action, and now it is assumed to be depending on Λ. Decompose the scalar field into $\phi = \psi + \chi$ such that $\psi \in C^\infty(\Sigma, M)[0, \Lambda']$ and $\chi \in C^\infty(\Sigma, M)(\Lambda', \Lambda]$, the path integral measure factorizes accordingly, i.e., $D\phi = D\psi\, D\chi$, so

$$e^{-S_{eff}[\Lambda,g]} = \int_{[0,\Lambda']} D\psi \int_{(\Lambda',\Lambda]} D\chi\ exp\left\{-\langle\psi + \chi, L(\Lambda, \psi + \chi, g)(\psi + \chi)\rangle\right\}$$

The action functional in the right hand side can be decomposed as

$$\begin{aligned}&\langle\psi+\chi, L(\Lambda,\psi+\chi,g)(\psi+\chi)\rangle\\&=\langle\psi, L(\Lambda,\psi,g)\psi\rangle+\langle\chi, L(\Lambda,\chi,g)\chi\rangle+S^{int}_{\Lambda}[\phi,\chi]\end{aligned}$$

where $S^{int}_{\Lambda}[\phi,\chi]$ is the effective interaction term. Since $S^{int}_{\Lambda}[\phi,\chi]$ must be at least cubic in the fields, we will ignore it in one-loop approximations. After droping higher order terms, we obtain the decomposition,

$$e^{-S_{eff}[\Lambda,g]}=\int_{[0,\Lambda']}D\psi exp\left\{-\langle\psi, L(\Lambda,\psi,g)\psi\rangle\right\}\int_{[\Lambda',\Lambda]}D\chi exp\left\{-\langle\chi, L(\Lambda,\chi,g)\chi\rangle\right\}$$

Denote the integrand using a new effective action $S(\psi,g)$,

$$e^{-S(\psi,g)}=exp\left\{-\langle\psi, L(\Lambda,\psi,g)\psi\rangle\right\}\int_{[\Lambda',\Lambda]}D\chi\, exp\left\{-\langle\chi, L(\Lambda,\chi,g)\chi\rangle\right\}\quad(7)$$

According to the scale invariance of RG flow, $S(\psi)$ is supposed to take the same form as before, namely

$$S(\psi,g)=\langle\psi, L(\Lambda',\psi,g)\psi\rangle\quad(8)$$

with a new energy scale Λ'. This assumption guarantees the effective action is invariant under the change of scales from Λ to Λ',

$$\begin{aligned}e^{-S_{eff}[\Lambda,g]}&=\textstyle\int_{[0,\Lambda]}D\phi\ exp\left\{-\langle\phi, L(\Lambda,\phi,g)\phi\rangle\right\}\\&=\textstyle\int_{[0,\Lambda']}D\psi\ exp\left\{-\langle\psi, L(\Lambda',\psi,g)\psi\rangle\right\}=e^{-S_{eff}[\Lambda',g]}\end{aligned}$$

Now under the scale invariance (8), we rewrite (7) as,

$$\langle\psi, L(\Lambda',\psi,g)\psi\rangle=\langle\psi, L(\Lambda,\psi,g)\psi\rangle-2\ln\int_{\Lambda'}^{\Lambda}D\chi\, exp\left\{-\frac{1}{2}\langle\chi, L(\Lambda,\chi,g)\chi\rangle\right\}$$

where a multiple $1/2$ is restored for the Gaussian integral, and the string tension T has been absorbed into the action. Suppose that $L(\Lambda,\psi,g)\psi=\lambda(\Lambda)\psi$ at the energy scale Λ for an eigenfunction ψ, and $L(\Lambda',\psi,g)\psi=\lambda(\Lambda')\psi$ is assumed to be the effective eigenvalue equation at Λ' for the same eigenfunction ψ. Note that the eigenfunction ψ can be chosen such that $\langle\psi,\psi\rangle=1$, and the definite integral is formally represented by the truncated Laplacian determinant, so we obtain

$$\lambda(\Lambda')=\lambda(\Lambda)+\ln\det(gD)|^{\Lambda}_{\Lambda'}$$

Using the heat kernel, it can be expressed as

$$\lambda(\Lambda') = \lambda(\Lambda) - \int_{\Lambda^{-2}}^{\Lambda'^{-2}} \frac{dt}{t} K(t, g, D) \tag{9}$$

where the integral variable is identified to be $t \sim \Lambda^{-2}$. Suppose that the energy level Λ is large and Λ' is very close to Λ so that $0 < \Lambda^{-2} < \Lambda'^{-2} < 1$, in this case the heat kernel is asymptotically expanded as usual

$$K(t, g, D) \sim \sum_{k \geq 0} t^{(k-n)/2} a_k(g, D)$$

where the multiple constant $(4\pi)^{-n/2}$ is absorbed in a_k.

Plug the asymptotic expansion into (9), we get

$$\lambda(\Lambda') = \lambda(\Lambda) - \sum_{k \geq 0} a_k(g, D) \int_{\Lambda^{-2}}^{\Lambda'^{-2}} t^{(k-n-2)/2} dt$$

There are three different parts in the above summation after we integrate it out. Here we introduce a new variable

$$\tau = \ln(\Lambda'^{-2}/\Lambda^{-2}), \quad i.e. \quad \Lambda'^{-2} = e^{\tau} \Lambda^{-2}$$

- $0 \leq k < n$, the first part is divergent if $\Lambda \to \infty$,

$$\sum_{0 \leq k < n} \frac{2a_k}{(k-n)} t^{(k-n)/2} |_{\Lambda^{-2}}^{\Lambda'^{-2}} = \sum_{0 \leq k < n} \frac{2a_k}{(k-n)} \Lambda^{n-k} (e^{-\tau(n-k)/2} - 1)$$

- $k > n$, this is the convergent part in the limit $\Lambda \to \infty$,

$$\sum_{k > n} \frac{2a_k}{(k-n)} t^{(k-n)/2} |_{\Lambda^{-2}}^{\Lambda'^{-2}} = \sum_{k > n} \frac{2a_k}{(k-n)} \frac{e^{\tau(k-n)/2} - 1}{\Lambda^{k-n}}$$

- $k = n$, this part has the expected logarithmic growth,

$$2 \ln(\Lambda/\Lambda')\, a_n(g, D)$$

Using the minimal subtraction renormalization scheme, we pick out this logarithmic growth term as the critical part, meanwhile we have to subtract the divergent part and ignore the convergent part. After the regularization

and renormalization, we get the expression of the effective eigenvalue at the energy scale Λ',

$$\lambda(\Lambda') = \lambda(\Lambda) - 2\ln(\Lambda/\Lambda')a_n(g, D)$$

On the right hand side, in addition to a renormalized eigenvalue $\lambda(\Lambda/\Lambda')$, a counterterm must be added to cancel the one loop singularity,

$$\lambda(\Lambda) = \lambda(\Lambda/\Lambda') + 2\ln(\Lambda/\Lambda')a_n(g, D)$$

Now the eigenvalue λ is physical and does not depend on the energy scale and the choice of renormalization schemes, that is, its variation with respect to $\tau = 2\ln(\Lambda/\Lambda')$ vanishes.

$$0 = \frac{\partial\lambda}{\partial\tau} = \frac{\partial}{\partial\tau}\lambda(\tau) + a_n(g, D)$$

In other words, the variation of the eigenvalue with respect to τ is

$$\frac{\partial}{\partial\tau}\lambda(\tau) = -a_n(g, D)$$

Finally, if we introduce a new parameter $t = -\tau$, then the right form under the RG flow is

$$\frac{\partial}{\partial t}\lambda(t) = a_n(g, D)$$

□

From [8], the regularized log-determinant can be defined as,

$$\ln\det(D, \Lambda) = -\zeta'(0, D) - \ln(\Lambda^2)\zeta(0, D) \tag{10}$$

for a positive elliptic second order differential operator D. From the proof of the above theorem, we know that

$$\ln\det(gD)|_{\Lambda'}^{\Lambda} = -2\ln(\Lambda/\Lambda')a_n(g, D) = -2\ln(\Lambda/\Lambda')\zeta(0, g, D)$$

In other words, if we set $g = 1$, then the variation of the log-determinant is,

$$\ln\det(D, \Lambda) - \ln\det(D, \Lambda') = -2\ln(\Lambda/\Lambda')\zeta(0, D)$$

which agrees with that derived from (10).

Therefore, under the RG flow, the log-determinant of the Polyakov action varies according to,

$$\frac{\partial}{\partial t}\ln\det(g,D,t) = a_n(g,D) = \zeta(0,g,D)$$

where $t = -\ln(\Lambda^2)$. From the next section, we know that the thermal entropy is proportional to the logarithm of the partition function,

$$S = \ln Z - \beta\frac{\partial}{\partial\beta}\ln Z \;\propto\; \ln Z$$

If we view the nonlinear sigma model (2) as a quantum statistical mechanical system, then the partition function is the trace of the heat kernel (3).

Corollary 1. *The variation of the thermal entropy of the Polyakov action is proportional to the top heat kernel coefficient along the RG flow,*

$$\frac{\partial S}{\partial t} \propto a_n(g,D) = \zeta(g,D) \tag{11}$$

Hence the variation of eigenvalues really reflects the variation of entropy under the RG flow. By the feature of heat kernel coefficients, we have to find a different formula for the variation of eigenvalues in odd dimensions.

4 A conjecture in odd dimensions

In this section, we will first go over the thermal entropy and its relation to the partition function and the free energy in statistical mechanics. After reviewing the irreversibility of the RG flow, i.e., the c, a, and F-theorems in conformal field theory, we will conjecture a formula for the variation of eigenvalues in odd dimensions based on the holographic principle.

4.1 Thermal entropy

Let us first recall some facts in classical and quantum statistical mechanics in the canonical ensemble, where the absolute temperature T is the principal thermodynamic variable and the total energy E may differ between microstates. As usual the inverse temperature is defined as $\beta = 1/kT$, where k is the Boltzmann constant, and as a convention one always sets $k = 1$.

In classical statistical mechanics, an ensemble is represented by a probability function P defined over the phase space. According to the Maxwell-Boltzmann distribution, the probability to find the system at the total energy E is given by

$$P = e^{\beta(F-E)}$$

where F is the Helmholtz free energy and is a constant for the ensemble. Or equivalently, the probability is expressed as,

$$P = \frac{1}{Z} e^{-\beta E} \quad \text{with} \quad Z = e^{-\beta F}$$

Z is called the canonical partition function. In other words, the free energy F can be used interchangeably with the partition function Z,

$$F = -\frac{1}{\beta} \ln Z$$

The Gibbs entropy is defined by the average of the logarithmic probability,

$$S = -\langle \ln P \rangle = \beta \langle E \rangle + \ln Z$$

where $\langle E \rangle$ denotes the average energy.

For convenience, one always expresses the thermodynamic variables in the partition function Z and its derivatives,

$$F = -\frac{1}{\beta} \ln Z, \quad \langle E \rangle = -\frac{\partial}{\partial \beta} \ln Z, \quad S = \ln Z - \beta \frac{\partial}{\partial \beta} \ln Z$$

The energy fluctuation is the variance of the energy

$$\sigma = \langle (E - \langle E \rangle)^2 \rangle = \frac{\partial^2}{\partial \beta^2} \ln Z \geq 0$$

and the derivative of the entropy is

$$\frac{\partial S}{\partial \beta} = -\beta \frac{\partial^2}{\partial \beta^2} \ln Z = -\beta \sigma \leq 0$$

To be complete, we also collect the thermodynamic variables expressed in the absolute temperature T and the partition function Z,

$$F = -T \ln Z, \quad \langle E \rangle = T^2 \frac{\partial \ln Z}{\partial T}, \quad S = \frac{\partial}{\partial T}(T \ln Z), \quad \frac{\partial S}{\partial T} = \frac{\sigma}{T^3} \geq 0$$

Notice that there is another interesting relation between the free energy F and the partition function Z,

$$S = -\frac{\partial F}{\partial T}$$

In addition, if the average energy vanishes, i.e., $\langle E \rangle = 0$, then

$$S = \ln Z = -\beta F$$

In quantum statistical mechanics, a statistical ensemble is represented by a density matrix ρ instead of the probability function,

$$\rho = e^{\beta(F-H)}$$

where H is the Hamiltonian of the canonical ensemble. By the normalization condition $\mathrm{Tr}\,\rho = 1$, one rewrites the above formula as,

$$e^{-\beta F} = \mathrm{Tr}\, e^{-\beta H}$$

Or equivalently, if $Z = \mathrm{Tr}\, e^{-\beta H}$ is called the partition function as usual, then

$$F = -\frac{1}{\beta} \ln Z$$

Now the von-Neumann entropy is defined as

$$S = -\mathrm{Tr}\, \rho \ln \rho = -\langle \ln \rho \rangle$$

From the relation between quantum statistical mechanics and quantum field theory, the partition function $Z_\beta = Tr(e^{-\beta H})$ will be effectively identified with the partition function $Z = \int d\phi exp\{\langle \phi, \Delta\phi \rangle\}$ from the path integral formalism for carefully chosen Hamiltonian H and Laplacian Δ. In this case, the inverse temperature β will be related to the energy scale by $\beta \sim \Lambda^{-2}$.

4.2 Holographic principle

In the study of renormalization group (RG) flow, the holographic principle is applied to understand the holographic RG flow in the AdS/CFT correspondence. In addition, entanglement entropy is used to probe the RG flow in the theory space of quantum field theories (QFTs), we know that entanglement

entropy is the same as thermal entropy for CFTs. In this section, we will conjecture a formula for the variation of eigenvalues in odd dimensions based on the holographic principle.

Ideally, one would like to construct a real-valued positive function $C(g, \Lambda)$, called the C-function, which depends on the coupling constant g and the energy scale Λ of an effective QFT. Let $C(g, \Lambda)$ vary along the RG flow, there are two desired properties: 1) $C(g, \Lambda)$ decreases monotonically from the UV to IR regime; 2) $C(g, \Lambda)$ is stationary (a constant independent of Λ) at the fixed points of the RG flow, i.e., at some conformal field theories (CFTs). The first milestone in this direction is Zamolodchikov's c-theorem in 2 dimensions [16]. More precisely, for 2d QFTs, there exists such a C-function so that it equals the central charge of the corresponding CFT at some fixed point of the RG flow. Furthermore, the irreversibility of the RG flow manifests itself in the relation,

$$c_{UV} > c_{IR}$$

Roughly speaking, the irreversibility of the RG flow means high energy theories have more information than theories at low energies. In essence, the irreversibility of the RG flow is equivalent to the 2nd law of thermaldynamics, that is, entropy must increase.

In even dimensions, there is a conformal anomaly or trace anomaly, that is, the trace of the energy-momentum tensor, i.e., T^{μ}_{μ}, is non-zero due to the conformal non-invariance after quantization. The conformal anomaly can be computed by one-loop approximation and zeta function regularization, in terms of heat kernel coefficients,

$$T^{\mu}_{\mu}(x) = a_n(x, D)$$

for a Laplace-type operator D.

In particular, the conformal anomaly in 2d can be locally expressed as,

$$T^{\mu}_{\mu}(x) = a_2(x, D) = \frac{1}{24\pi} R$$

While in a 2d CFT, it is usually written as,

$$\langle T^{\mu}_{\mu} \rangle = -\frac{c}{24\pi} R$$

where c is the central charge. The irreversibility of the RG flow, i.e., $c_{UV} > c_{IR}$, was proved based on computations involving the energy-momentum tensor. Later it was also interpreted as the entanglement entropy of an interval, since the central charge also appears in the von-Neumann entropy.

Next in 4d, one has the local form of the conformal anomaly,

$$T^{\mu}_{\mu}(x) = a_4(x, D) = \frac{1}{2880\pi^2}(R^{ijkl}R_{ijkl} - R^{ij}R_{ij} - 30\Delta R + 5R^2)$$

In CFT, it is alwayls written as

$$\langle T^{\mu}_{\mu} \rangle = -\frac{1}{16\pi^2}(aE_4 - cW^2)$$

where E_4 is the Euler density and W^2 is the Weyl tensor squared,

$$E_4 = R^2_{\mu\nu\rho\sigma} - 4R^2_{\mu\nu} + R^2, \quad W^2_{\mu\nu\rho\sigma} = R^2_{\mu\nu\rho\sigma} - 2R^2_{\mu\nu} + \frac{1}{3}R^2$$

Again, a and c are two central charges in CFT. For conformally flat spaces such as the four-sphere, one has $W^2 = 0$, so in this case the conformal anomaly is only depending on the central charge a,

$$\langle T^{\mu}_{\mu} \rangle = -\frac{a}{16\pi^2}E_4$$

As an analogy of the c-theorem in 2d, there is a so-called a-theorem in 4d [9], that is, along the RG flow,

$$a_{UV} > a_{IR}$$

There is no conformal anomaly or trace anomaly in odd dimensions. Under the holographic RG flow, an F-theorem was proved in 3d [3], that is, along the RG flow,

$$F_{UV} > F_{IR}$$

where F is the free energy of the three sphere,

$$F = -\frac{1}{\beta}\ln Z_{S^3} = -T \ln Z_{S^3}$$

Later F was also interpreted as the (renormalized) entanglement entropy in [12].

Proposition 1. *Under the holographic principle, the free energy changes along the RG flow as*

$$\frac{\partial F}{\partial t} = \frac{1}{2}\zeta'(0, D) - t\zeta(0, D) = \frac{1}{2}\zeta'(0, D) - ta_{2k}(D) \tag{12}$$

Proof. Under the AdS/CFT correspondence, we consider the bulk-boundary duality. We assume the bulk geometry is given by a $2k$-dimensional sphere S^{2k}, i.e., the Euclidean version of the AdS_{2k} space, and the boundary field theory is described by a $(2k-1)$-dimensional CFT. The basic assumption of the holographic principle is that the partition functions are the same

$$Z_{CFT} = Z_{S^{2k}}$$

So the free energy of the odd dimensional boundary theory is

$$F = -T \ln Z_{CFT} = -T \ln Z_{S^{2k}} = -\tau \ln Z_{S^{2k}}$$

where the absolute temperature T is identified with the parameter $\tau = \ln \Lambda^2$ in the RG flow since T has units as in energy. For scalar fields, the bulk partition function is given by the one-loop approximation,

$$\ln Z_{S^{2k}} = -\frac{1}{2} \ln \det D = \frac{1}{2}[\zeta'(0, D) + \tau\zeta(0, D)]$$

where we use the regularized log-determinant (10) of the bulk Laplacian D. So the free energy is written as,

$$F = -\frac{1}{2}[\tau\zeta'(0, D) + \tau^2\zeta(0, D)]$$

Hence the derivative with respect to τ is

$$\frac{\partial F}{\partial \tau} = -\frac{1}{2}\zeta'(0, D) - \tau\zeta(0, D)$$

If we introduce a new parameter $t = -\tau$, then the right form of the variation of F along the RG flow is given by

$$\frac{\partial F}{\partial t} = \frac{1}{2}\zeta'(0, D) - t\zeta(0, D)$$

□

If the thermal entropy S is proportional to the minus free energy $-F$ in odd dimensions,

$$S \propto -F$$

then the variation of entropy along the RG flow is

$$\frac{\partial S}{\partial t} \propto -\frac{1}{2}\zeta'(0,D) + t\zeta(0,D)$$

Next we consider some eigenvalue representing the entropy or the minus free energy in odd dimensions, we have the following conjecture.

Conjecture 1. *If the holographic Laplacian in the boundary theory is assumed to be $\hat{D}$, which is dual to the bulk Laplacian D, then the variation of eigenvalues of $\hat{D}$ along the RG flow in $(2k-1)$ dimensions is*

$$\frac{\partial \lambda}{\partial t} = -\frac{1}{2}\zeta'(0,D) + ta_{2k}(D) \tag{13}$$

In particular, if we only approximate the above formula to the first order, then the variation of eigenvalues is

$$\frac{\partial \lambda}{\partial t} = -\frac{1}{2}\zeta'(0,D) \tag{14}$$

5 Discussion

The irreversibility of the renormalization group (RG) flow is closely related to the monotonicity of entropy. And the fixed points of the RG flow, i.e., the corresponding conformal field theories (CFTs), play an important role in understanding such irreversibility. The eigenvalues derived from entropy functionals inherit the monotonicity of entropy. From the one-loop approximation of a quantum field theory (QFT), the effective partition function can be computed based on heat kernel expansion and zeta function regularization. However, a general heat kernel coefficient does not have the monotonicity, neither do the eigenvalues. In order to establish the monotonicity, one has to decompose a heat kernel coefficient into different parts. For example, it is possible to find the monotonicity from the Euler density in 4d.

In odd dimensions, there is no conformal anomaly, so it is only possible to find the monotonicity based on the holographic principle. We have learned something interesting from the F-theorem in 3d. The free energy in odd

dimensions plays the same role as the partition function in even dimensions. There are too many mysterious things that we do not understand in the holographic principle, so we end this paper with a conjecture. We conjectured a formula for the variation of eigenvalues in odd dimensions.

In this paper, we are mainly interested in the Polyakov action (2) and the corresponding Laplacian (5). It is straightforward to generalize the Polyakov action to include a dilaton field, and it may take more effects to include B-fields and other fields in string theory. It is also interesting to apply a similar strategy to study spinor fields or gauge fields evolving along the RG flow.

We also hope to generalize this work into the noncommutative world. In noncommutative geometry, a Dirac operator in a spectral triple plays the role of a metric in Riemannian geometry. Instead of deformations of a Dirac operator, it is easier to consider deformations of eigenvalues of the Dirac Laplacian under some geometric flow or the RG flow. There are a lot of things one can try to study the conformal structure in noncommutative geometry.

References

[1] X.D. Cao. First eigenvalues of geometric operators under the Ricci flow. *Proc. Amer. Math. Soc.*, 136(11):4075–4078, 2008.

[2] M. Carfora. Renormalization group and the Ricci flow. *Milan J. Math.*, 78:319–353, 2010.

[3] H. Casini, M. Huerta, and R. C. Myers. Towards a derivation of holographic entanglement entropy. *JHEP*, 05:36, 2011.

[4] L.F. Di Cerbo. Eigenvalues of the laplacian under the Ricci flow. *Rendiconti di Matematica*, 27:183–195, 2007.

[5] D. Friedan. Nonlinear models in $2 + \epsilon$ dimensions. *Phys.Rev.Lett.*, 45:1057, 1980.

[6] D. Friedan. Nonlinear models in $2 + \epsilon$ dimensions. *Ann. of Phys.*, 163:318–419, 1985.

[7] K. Gawedzki. Lectures on conformal field theory. In P. Deligne, editor, *Quantum fields and strings: a course for mathematicians*, pages 727–805. AMS, 2000.

[8] S.W. Hawking. Zeta function regularization of path integrals in curved spacetime. *Comm. Math. Phys.*, 55(2):133–148, 1977.

[9] Z. Komargodski and A. Schwimmer. On renormalization group flows in four dimensions. *JHEP*, 12:99, 2011.

[10] J.F. Li. Eigenvalues and energy functionals with monotonicity formulae under Ricci flow. *Math. Ann.*, 338(4):927–946, 2007.

[11] X.D. Li. From the Boltzmann H-theorem to Perelman's W-entropy formula for the Ricci flow. In H. Chen, Y.M. Long, and Y. Nishiura, editors, *Emerging Topics on Differential Equations and Their Applications*, pages 68–74. World Scientific, 2012.

[12] H. Liu and M. Mezei. A refinement of entanglement entropy and the number of degrees of freedom. *JHEP*, 04:162, 2013.

[13] Li Ma. Eigenvalue monotonicity for the Ricci-Hamilton flow. *Ann. Global Anal. Geom.*, 29(3):287–292, 2006.

[14] G. Perelman. The entropy formula for the Ricci flow and its geometric applications. 2002. arXiv: 0211159.

[15] D.V. Vassilevich. Heat kernel expansion: User's manual. *Phys.Rept.*, 388:279–360, 2003.

[16] A. Zamolodchikov. Irreversibility of the flux of the renormalization group in a 2D field theory. *JETP Lett.*, 43:730–732, 1986.